

### Anti-3-Nitrotyrosine antibody [39B6] ab61392

★★★★★ **9 Abreviews**   **130 References**   画像数 6

#### 製品の概要

製品名	Anti-3-Nitrotyrosine antibody [39B6]
製品の詳細	Mouse monoclonal [39B6] to 3-Nitrotyrosine
由来種	Mouse
特異性	No detectable crossreactivity with non-nitrated tyrosine. Not species specific.
アプリケーション	<b>適用あり:</b> IHC-P, WB
種交差性	<b>交差種:</b> Species independent
免疫原	Full length protein corresponding to 3-Nitrotyrosine. 3-(4-hydroxy-3-nitrophenylacetamido) propionic acid-bovine serum albumin
ポジティブ・コントロール	SIN-1 (3-morpholinosydnonimine) treated BSA. Use between 1-2.5mM SIN-1 for 1-24 hours, grow cells under normal growth conditions. Please see Yasuda S, Idell S, Liu MC. Biochem J. 2007 Jan 15;401(2):497-503. PMID: 17002600. IHC-P: Mouse back skin and inflamed colon tissue; Human colon carcinoma tissue. Rat liver tissue. WB: A549 cell lysate.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.09% Sodium azide Constituents: PBS, 50% Glycerol
精製度	Protein G purified
ポリ/モノ	モノクローナル
クローン名	39B6

アイソタイプ IgG2a  
軽鎖の種類 kappa

## アプリケーション

**The Abpromise guarantee** Abpromise保証は、次のテスト済みアプリケーションにおけるab61392の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
IHC-P	★★★★★ (3)	1/100.
WB	★★★★★ (6)	1/1000. We would recommend a 1:1000 dilution for the antibody, for 2 hours at RT. For blocking, use milk.

## ターゲット情報

**関連性**


Protein tyrosine nitration results in a post-translational modification that is increasingly receiving attention as an important component of nitric oxide signaling. While multiple nonenzymatic mechanisms are known to be capable of producing nitrated tyrosine residues, most tyrosine nitration events involve catalysis by metalloproteins such as myeloperoxidase, eosinophilperoxidase, myoglobin, the cytochrome P-450s, superoxide dismutase and prostacyclin synthase. Various studies have shown that protein tyrosinenitration is limited to specific proteins and that the process is selective. For example, exposure of human surfactant protein A, SP-A, to oxygen-nitrogen intermediates generated by activated alveolar macrophages resulted in specific nitration of SP-A at tyrosines 164 and 166, while addition of 1.2 mMCO 2 resulted in additional nitration at tyrosine 161. The presence of nitrotyrosine-containing proteins has shown high correlation to disease states such as atherosclerosis, Alzheimer's disease, Parkinson's disease and amyotrophic lateral sclerosis.

## 画像



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-3-Nitrotyrosine antibody [39B6] (ab61392)


ab61392 staining 3-nitrotyrosine in Bouin's fixed paraffin-embedded Rat liver tissue at 1/1000 dilution. FITC-conjugated goat anti-mouse IgG(green) was used as a secondary antibody.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-3-Nitrotyrosine antibody [39B6] (ab61392)

ab61392 staining 3-nitrotyrosine in Bouin's fixed paraffin-embedded backskin sections of transgenic mice.

Blocker: 5% goat serum in buffer. Buffer: 99% washing buffer (7g BSA and 222ul gelatin topped up to 1L dH<sub>2</sub>O), 1% heat inactivated goat serum. Primary Dilution: 1/100 for 1 hour at room temperature. Secondary: FITC-conjugated goat anti-mouse IgG, 1/50 for 1 hour at room temperature.



Western blot - Anti-3-Nitrotyrosine antibody [39B6] (ab61392)

This image is courtesy of an anonymous Abreview

**All lanes :** Anti-3-Nitrotyrosine antibody [39B6] (ab61392) at 1/3000 dilution

**All lanes :** Mouse skeletal muscle

Lysates/proteins at 20 µg per lane.

#### Secondary


**All lanes :** Goat anti-Mouse HRP conjugated at 1/6000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

**Exposure time:** 4 seconds


Blocked with 5% milk for 2 hours.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-3-Nitrotyrosine antibody [39B6] (ab61392)

Formalin-fixed, paraffin-embedded mouse inflamed colon tissue stained for Nitrotyrosine using ab61392 at 1/1000000 dilution in immunohistochemical analysis.


Secondary Antibody: Biotin Goat Anti-Mouse at 1/2000 for 1 hour at RT. Counterstain: Mayer Hematoxylin (purple/blue) nuclear stain at 200 µl for 2 minutes at RT.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-3-Nitrotyrosine antibody [39B6] (ab61392)

Formalin-fixed, paraffin-embedded human colon carcinoma tissue stained for Nitrotyrosine using ab61392 at 1/25000 dilution in immunohistochemical analysis.

Secondary Antibody: Biotin Goat Anti-Mouse at 1/2000 for 1 hour at RT. Counterstain: Mayer Hematoxylin (purple/blue) nuclear stain at 200 µl for 2 minutes at RT.



Western blot - Anti-3-Nitrotyrosine antibody [39B6] (ab61392)

Anti-3-Nitrotyrosine antibody [39B6] (ab61392) at 1/1000 dilution (2.5 hours at RT with shaking) + A549 (Human lung carcinoma cell line) cell lysate at 15 µg

#### Secondary

Goat anti-mouse IgG HRP, 1 hour at RT with shaking at 1/1000 dilution

Block: 5% Skim milk powder in TBST.

Color Development: Chemiluminescent for HRP for 5 mins at RT.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

**Our Abpromise to you: Quality guaranteed and expert technical support**

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors